arXiv:1211.5828v1 [math.AG] 26 Nov 2012

A GOING-UP THEOREM

YING ZONG

We prove a generalization of Zariski's irregularity theorem on cyclic multiple planes [1], p. 508.

1. Notations

Let p be a fixed prime number, $\Lambda = \mathbf{Z}/p\mathbf{Z}$.

Given a scheme X, and $* \in \{+,-,b,\emptyset\}$, $D^*(X,\Lambda)$ denotes the derived category of the category of sheaves of Λ-modules on X for the étale topology.

If $T : D^*(X,\Lambda) \to A$ is a cohomological functor with values in an abelian category, $L \in \mathrm{Ob}\ D^*(X,\Lambda)$, $i \in \mathbf{Z}$, let $T^i(L)$ denote $T(L[i])$.

2. A Going-up theorem

Theorem. — *Let X be a scheme, $i : D \hookrightarrow X$ a closed immersion, U the complement of D in X. Suppose U is locally connected. Suppose $\pi : Y \to X$ is an X-scheme, entier, surjective over X, radical over D, and finite, étale, Galois over U of group a p-group. If $T : D^*(X,\Lambda) \to A$ is a cohomological functor with values in an abelian category, K is an object of $D^*(X,\Lambda)$ such that $T(K) \to T(i_*i^*K)$ is surjective and $T^1(K) = 0$, then $T^1(\pi_*\pi^*K) = 0$.*

Remarks. — i) In particular, if $H^0(X,\Lambda) \to H^0(D,\Lambda)$ is surjective, and $H^1(X,\Lambda) = 0$, then $H^1(Y,\Lambda) = 0$. For D a connected curve on the complex projective plane X, this is essentially Zariski's theorem, *loc.cit.*

ii) Write $j : U \hookrightarrow X$, $j' : \pi^{-1}(U) \hookrightarrow Y$ for the open immersions. Let $T = H^0R\Gamma : D^+(X,\Lambda) \to (Ab)$, $n \in \mathbf{Z}$, $K = j_!\Lambda[n-1]$. Since $i^*K = 0$, one finds that $H^n(Y, j'_!\Lambda) = 0$, if $H^n(X, j_!\Lambda) = 0$.

iii) The following analogue is proven in the same way :

Let X be a topological space, $i : D \hookrightarrow X$ a closed subspace, U the complement of D in X. Suppose U is locally connected and locally path simply connected. Suppose $\pi : Y \to X$ is an X-space, proper, surjective, with finite fibres, $\pi^{-1}(D) \to D$ is bijective, $\pi^{-1}(U) \to U$ is a Galois covering of Galois group a p-group. If $T : D^*(X,\Lambda) \to A$ is a *cohomological functor with values in an abelian category, K is an object of $D^*(X, \Lambda)$ such that $T(K) \to T(i_*i^*K)$ is surjective and $T^1(K) = 0$, then $T^1(\pi_*\pi^*K) = 0$.*

Proof of the theorem. — Write $j : U \hookrightarrow X$ for the open immersion.

Every object L of $D^*(X, \Lambda)$ sits in the localization triangle (SGA 4, Éxposé XVII, 5.1.16)

$$j_!j^*L \to L \to i_*i^*L.$$

Its T-cohomology sequence reads as

$$T^1(j_!j^*L) \to T^1(L) \to T^1(i_*i^*L) \xrightarrow{\delta_L} T^2(j_!j^*L).$$

If, for $L = \pi_*\pi^*K$, one has that $T^1(j_!j^*L) = 0$ and

$$\delta_L : T^1(i_*i^*L) \to T^2(j_!j^*L)$$

is injective, then $T^1(L) = 0$.

Prove $T^1(j_!j^*\pi_*\pi^*K) = 0$:

The morphism $\pi^{-1}(U) \to U$ being finite, étale, Galois of group a p-group G, the Λ-module $j^*\pi_*\Lambda$ is locally constant, constructible, and at any geometric point $\overline{u}$ of U, the representation

$$\rho_{\overline{u}} : \pi_1(U, \overline{u}) \to \mathrm{GL}((j^*\pi_*\Lambda)_{\overline{u}})$$

factors through G. Consequently, the monodromy $\mathrm{Im}(\rho_{\overline{u}})$ is unipotent, and

$$F^1 := \mathrm{Coker}(\Lambda \to j^*\pi_*\Lambda)$$

admits a finite filtration F^n by sub-Λ-modules such that each successive quotient F^n/F^{n+1} is isomorphic to Λ.

The exact sequences

$$0 \to \Lambda \to j^*\pi_*\Lambda \to F^1 \to 0$$

$$0 \to F^{n+1} \to F^n \to \Lambda \to 0$$

applied by

$$? \ \mapsto \ j^*K \overset{\mathbf{L}}{\otimes}_\Lambda \ ?$$

give rise to triangles

$$j^*K \to j^*K \overset{\mathbf{L}}{\otimes}_\Lambda j^*\pi_*\Lambda \to j^*K \overset{\mathbf{L}}{\otimes}_\Lambda F^1$$

$$j^*K \overset{\mathbf{L}}{\otimes}_\Lambda F^{n+1} \to j^*K \overset{\mathbf{L}}{\otimes}_\Lambda F^n \to j^*K.$$

Further applied by the functor $Tj_!$, the triangles induce cohomology sequences

$$(*)\quad T^1(j_!j^*K) \to T^1(j_!(j^*K \overset{\mathbf{L}}{\otimes}_\Lambda j^*\pi_*\Lambda)) \to T^1(j_!(j^*K \overset{\mathbf{L}}{\otimes}_\Lambda F^1))$$

$$(*_n)\quad T^1(j_!(j^*K \overset{\mathbf{L}}{\otimes}_\Lambda F^{n+1})) \to T^1(j_!(j^*K \overset{\mathbf{L}}{\otimes}_\Lambda F^n)) \to T^1(j_!j^*K).$$

Observe that

$$T^1(j_!j^*K) = 0$$

using the exact sequence

$$T(K) \to T(i_*i^*K) \xrightarrow{\delta_K} T^1(j_!j^*K) \to T^1(K)$$

and the assumption that $T(K) \to T(i_*i^*K)$ is surjective and $T^1(K) = 0$.

By descending induction on n, $(*_n)$ and $(*)$ imply that

$$(**_n)\qquad T^1(j_!(j^*K \overset{\mathbf{L}}{\otimes}_\Lambda F^n)) = 0, \quad \forall\ n \in \mathbf{Z}$$

$$(**)\qquad T^1(j_!(j^*K \overset{\mathbf{L}}{\otimes}_\Lambda j^*\pi_*\Lambda)) = 0.$$

Then, the projection formula

$$\pi_*\pi^*K = K \overset{\mathbf{L}}{\otimes}_\Lambda \pi_*\Lambda$$

(SGA 4, Éxposé XVII, Proposition 5.2.9) concludes the vanishing

$$T^1(j_!j^*\pi_*\pi^*K) = T^1(j_!j^*(K \overset{\mathbf{L}}{\otimes}_\Lambda \pi_*\Lambda)) = T^1(j_!(j^*K \overset{\mathbf{L}}{\otimes}_\Lambda j^*\pi_*\Lambda)) \overset{(**)}{=} 0.$$

Next, show

$$\delta_L : T^1(i_*i^*\pi_*\pi^*K) \to T^2(j_!j^*\pi_*\pi^*K)$$

is injective :

Let $i' : \pi^{-1}(D) \hookrightarrow Y$ denote the closed immersion, $\pi' : \pi^{-1}(D) \to D$ the restriction of π to $\pi^{-1}(D)$.

The theorem of base change (SGA 4, Éxposé VIII, Proposition 5.5)

$$i^*\pi_* = \pi'_*(i')^*$$

gives the equality

$$T^1(i_*i^*\pi_*\pi^*K) = T^1(i_*\pi'_*(i')^*\pi^*K) = T^1(i_*\pi'_*(\pi')^*i^*K).$$

And the adjunction morphism

$$i^*K \to \pi'_*(\pi')^*i^*K$$

is an isomorphism, for $\pi' : \pi^{-1}(D) \to D$ is entier, radical, surjective (SGA 4, Éxposé VIII, Théorème 1.1).

Hence, canonically,

$$T^1(i_*i^*\pi_*\pi^*K) = T^1(i_*i^*K).$$

The homomorphism

$$\delta_L : T^1(i_*i^*\pi_*\pi^*K) \to T^2(j_!j^*\pi_*\pi^*K)$$

factors, up to the identification

$$T^1(i_*i^*\pi_*\pi^*K) = T^1(i_*i^*K),$$

as

$$\delta_L : T^1(i_*i^*K) \xrightarrow{\delta_K} T^2(j_!j^*K) \to T^2(j_!j^*\pi_*\pi^*K).$$

To show

$$\delta_L : T^1(i_*i^*\pi_*\pi^*K) \to T^2(j_!j^*\pi_*\pi^*K)$$

is injective, it suffices to show

$$\delta_K : T^1(i_*i^*K) \to T^2(j_!j^*K)$$

$$T^2(j_!j^*K) \to T^2(j_!j^*\pi_*\pi^*K)$$

are both injective.

That $\delta_K : T^1(i_*i^*K) \to T^2(j_!j^*K)$ is injective is immediate from the exact sequence

$$T^1(K) \to T^1(i_*i^*K) \xrightarrow{\delta_K} T^2(j_!j^*K)$$

and that $T^1(K) = 0$.

Finally, $T^2(j_!j^*K) \to T^2(j_!j^*\pi_*\pi^*K)$ is injective :

This follows from the triangle

$$j^*K \to j^*K \overset{\mathbf{L}}{\otimes}_\Lambda j^*\pi_*\Lambda \to j^*K \overset{\mathbf{L}}{\otimes}_\Lambda F^1,$$

its associated cohomology sequence

$$T^1(j_!(j^*K \overset{\mathbf{L}}{\otimes}_\Lambda F^1)) \to T^2(j_!j^*K) \to T^2(j_!(j^*K \overset{\mathbf{L}}{\otimes}_\Lambda j^*\pi_*\Lambda)),$$

the vanishing

$$(**_1) \quad T^1(j_!(j^*K \overset{\mathbf{L}}{\otimes}_\Lambda F^1)) = 0,$$

and the identity

$$j_!j^*\pi_*\pi^*K = j_!(j^*K \overset{\mathbf{L}}{\otimes}_\Lambda j^*\pi_*\Lambda).$$

REFERENCES

1. O. Zariski. On the irregularity of cyclic multiple planes. *Annals of Mathematics*, 32, 1931.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TORONTO
E-mail address: `zongying@math.toronto.edu`